

1 Introduction

Intrinsic spin and orbital angular momentum are fundamentally the same. One difference between them appears to be that while orbital angular momentum has a wave-mechanical description through Legendre's equation, spin does not. However, in this paper, we propose a differential eigenvalue equation for spin 1/2.

The arguments leading to our proposed equation are based on the Landé interpretation of quantum mechanics[1-4]. In this interpretation, the probability-amplitude character of an eigenfunction or wave function is stressed. Such a probability amplitude connects well-defined initial and final states[5]. Perhaps the most important such probability amplitudes result from solution of the time-independent Schrödinger equation

$$-\frac{\hbar^2}{2m}\nabla^2\psi(\mathbf{r}) + V(\mathbf{r})\psi(\mathbf{r}) = E\psi(\mathbf{r}). \quad (1)$$

The eigenfunction $\psi(\mathbf{r})$ connects an initial state defined by the eigenvalue E and a final state defined by the eigenvalue $\mathbf{r}$. To emphasize the initial and final state, we write the eigenfunction as $\psi(\mathbf{r}) = \psi(E_n; \mathbf{r})$, so that the equation reads

$$-\frac{\hbar^2}{2m}\nabla^2\psi(E_n; \mathbf{r}) + V(\mathbf{r})\psi(E_n; \mathbf{r}) = E_n\psi(E_n; \mathbf{r}). \quad (2)$$

The Schrödinger equation is, of course, a typical eigenvalue equation. Another example is the Legendre equation

$$L^2(\theta, \varphi)Y_{lm}(\theta, \varphi) = l(l+1)\hbar^2 Y_{lm}(\theta, \varphi). \quad (3)$$

In this equation, the initial state corresponds to the eigenvalues $l(l+1)\hbar^2$ and $m\hbar$ while the final state corresponds to the angular position (θ, φ) [6]. For this reason, we may write the eigenfunctions as $Y_{lm}(\theta, \varphi) = Y(l, m; \theta, \varphi)$.

We notice that in both the Schrödinger and the Legendre equations, the operator is cast in terms of the variables which also define the final eigenvalue. We take this to be a general feature, which we exploit in our search for a differential equation for spin.

2 Spin Eigenvectors

As conventionally treated, spin is such that both the initial and final states correspond to discrete eigenvalues. A differential eigenvalue equation requires a continuous variable for the operator to act on. But, the Pauli spin vectors

$$[\xi_{z,z}^+] = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{and} \quad [\xi_{z,z}^-] = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (4)$$

do not appear to offer a continuous variable which might make possible a differential treatment of spin. However, the Pauli vectors are specialised forms of the generalized spin eigenvectors[5]

$$[\xi_{\hat{\mathbf{a}},\hat{\mathbf{b}}}^+] = \begin{pmatrix} \chi((+\frac{1}{2})^{(\hat{\mathbf{a}})}; (+\frac{1}{2})^{(\hat{\mathbf{b}})}) \\ \chi((+\frac{1}{2})^{(\hat{\mathbf{a}})}; (-\frac{1}{2})^{(\hat{\mathbf{b}})}) \end{pmatrix} = \begin{pmatrix} \cos \theta/2 \cos \theta'/2 + e^{i(\varphi-\varphi')} \sin \theta/2 \sin \theta'/2 \\ \cos \theta/2 \sin \theta'/2 - e^{i(\varphi-\varphi')} \sin \theta/2 \cos \theta'/2 \end{pmatrix} \quad (5)$$

and

$$[\xi_{\hat{\mathbf{a}},\hat{\mathbf{b}}}^-] = \begin{pmatrix} \chi((-\frac{1}{2})^{(\hat{\mathbf{a}})}; (+\frac{1}{2})^{(\hat{\mathbf{b}})}) \\ \chi((-\frac{1}{2})^{(\hat{\mathbf{a}})}; (-\frac{1}{2})^{(\hat{\mathbf{b}})}) \end{pmatrix} = \begin{pmatrix} \sin \theta/2 \cos \theta'/2 - e^{i(\varphi-\varphi')} \cos \theta/2 \sin \theta'/2 \\ \sin \theta/2 \sin \theta'/2 + e^{i(\varphi-\varphi')} \cos \theta/2 \cos \theta'/2 \end{pmatrix}. \quad (6)$$

They result when we set all the angles equal to zero.

The elements of the eigenvectors Eqns. (5) and (6) are probability amplitudes $\chi((m_i)^{(\hat{\mathbf{a}})}; (m_f)^{(\hat{\mathbf{b}})})$ for spin projection measurements from the direction $\hat{\mathbf{a}}$ (polar angles (θ', φ')) to the direction $\hat{\mathbf{b}}$ (polar angles (θ, φ))[5]. The projection quantum numbers m_i and m_f give the spin projections corresponding to the appropriate directions. This probability amplitude represents the probability of obtaining the spin projection $m_f \hbar$ along the direction $\hat{\mathbf{b}}$ if the spin projection is initially $m_i \hbar$ along the direction $\hat{\mathbf{a}}$. Our notation for the eigenvectors takes cognisance of the quantization directions that define the probability amplitudes that constitute the elements of the eigenvectors. This is why the Pauli spin vectors Eqns. (4) contain the indices z and z . They contain as their elements probability amplitudes for spin projection measurements from the z direction to the z direction.

The most important aspect of the eigenvectors Eqns. (5) and (6) is that the polar angles (θ, φ) of the final quantization direction $\hat{\mathbf{b}}$ give us continuous

variables on which a differential operator may act. In a preliminary search for a differential operator for spin, we may simplify our task by setting the angles referring to the initial eigenvalue to fixed values, so that we have only the angles for the final observable as continuous variables. It is easiest to set the polar angles of $\hat{\mathbf{a}}$ to $(\theta', \varphi') = (0, 0)$, so that $\hat{\mathbf{a}}$ defines the z axis. In that case, we obtain

$$[\xi_{z,\hat{\mathbf{b}}}^+] = \begin{pmatrix} \chi((+\frac{1}{2})^{(\hat{\mathbf{k}})}; (+\frac{1}{2})^{(\hat{\mathbf{b}})}) \\ \chi((+\frac{1}{2})^{(\hat{\mathbf{k}})}; (-\frac{1}{2})^{(\hat{\mathbf{b}})}) \end{pmatrix} = \begin{pmatrix} \cos \theta/2 \\ -e^{i\varphi} \sin \theta/2 \end{pmatrix} \quad (7)$$

and

$$[\xi_{z,\hat{\mathbf{b}}}^-] = \begin{pmatrix} \chi((-\frac{1}{2})^{(\hat{\mathbf{k}})}; (+\frac{1}{2})^{(\hat{\mathbf{b}})}) \\ \chi((-\frac{1}{2})^{(\hat{\mathbf{k}})}; (-\frac{1}{2})^{(\hat{\mathbf{b}})}) \end{pmatrix} = \begin{pmatrix} \sin \theta/2 \\ e^{i\varphi} \cos \theta/2 \end{pmatrix}, \quad (8)$$

which we write in the more symmetrical forms

$$[\xi_{z,\hat{\mathbf{b}}}^+] = \begin{pmatrix} e^{-i\varphi/2} \cos \theta/2 \\ -e^{i\varphi/2} \sin \theta/2 \end{pmatrix} \quad \text{and} \quad [\xi_{z,\hat{\mathbf{b}}}^-] = \begin{pmatrix} e^{-i\varphi/2} \sin \theta/2 \\ e^{i\varphi/2} \cos \theta/2 \end{pmatrix}. \quad (9)$$

3 Differential Operators

We seek a differential operator $[S_{z,\hat{\mathbf{b}}}]$ such that

$$[S_{z,\hat{\mathbf{b}}}][\xi_{z,\hat{\mathbf{b}}}^\pm] = \pm \frac{1}{2} \hbar [\xi_{z,\hat{\mathbf{b}}}^\pm]. \quad (10)$$

The notation for the operator again takes into account the initial and final quantization directions contained in the eigenvectors of the operator. Since the solutions are 2×1 matrices, of which there are two independent ones, the operator must be a 2×2 matrix, with elements that are differential operators. We therefore adopt the form

$$[S_{z,\hat{\mathbf{b}}}] = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad (11)$$

for the operator. As the operator is analogous to L_z in the theory of orbital angular momentum, we assume that the differential operators are of first degree in the variables. Therefore, the elements are taken to have the structure

$$A = a_1 \frac{\partial}{\partial \theta} + a_2 \frac{\partial}{\partial \varphi}, \quad (12)$$

$$B = b_1 \frac{\partial}{\partial \theta} + b_2 \frac{\partial}{\partial \varphi}, \quad (13)$$

$$C = c_1 \frac{\partial}{\partial \theta} + c_2 \frac{\partial}{\partial \varphi} \quad (14)$$

and

$$D = d_1 \frac{\partial}{\partial \theta} + d_2 \frac{\partial}{\partial \varphi}, \quad (15)$$

with $a_1, a_2, \dots, d_2$ to be determined from the condition Eqn. (10).

By substituting into the eigenvalue equation, we find

$$b_1 = b_2 = c_1 = c_2 = 0, \quad (16)$$

$$a_1 = -\hbar \sin \theta, \quad a_2 = i\hbar \cos \theta \quad (17)$$

and

$$d_1 = \hbar \sin \theta, \quad d_2 = i\hbar \cos \theta. \quad (18)$$

Hence

$$[S_{z, \hat{\mathbf{b}}}] = \hbar \begin{pmatrix} -\sin \theta \frac{\partial}{\partial \theta} + i \cos \theta \frac{\partial}{\partial \varphi} & 0 \\ 0 & \sin \theta \frac{\partial}{\partial \theta} + i \cos \theta \frac{\partial}{\partial \varphi} \end{pmatrix}. \quad (19)$$

Coming now to the x component, we observe that the eigenvectors for the x component of the spin are[7]

$$[\xi_{x, \hat{\mathbf{b}}}^+] = \frac{1}{\sqrt{2}} \begin{pmatrix} (\sin \theta/2 + \cos \theta/2)e^{-i\varphi/2} \\ (\cos \theta/2 - \sin \theta/2)e^{i\varphi/2} \end{pmatrix} \quad (20)$$

$$[\xi_{x, \hat{\mathbf{b}}}^-] = \frac{1}{\sqrt{2}} \begin{pmatrix} (\sin \theta/2 - \cos \theta/2)e^{-i\varphi/2} \\ (\cos \theta/2 + \sin \theta/2)e^{i\varphi/2} \end{pmatrix} \quad (21)$$

Assuming for the operator $[S_{x, \hat{\mathbf{b}}}]$ the form

$$[S_{x, \hat{\mathbf{b}}}] = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix}, \quad (22)$$

with

$$A = a_1 \frac{\partial}{\partial \theta} + a_2 \frac{\partial}{\partial \varphi} \quad (23)$$

and

$$D = d_1 \frac{\partial}{\partial \theta} + d_2 \frac{\partial}{\partial \varphi}, \quad (24)$$

we find

$$[S_{x,\hat{\mathbf{b}}}] = \hbar \begin{pmatrix} \cos \theta \frac{\partial}{\partial \theta} + i \sin \theta \frac{\partial}{\partial \varphi} & 0 \\ 0 & -\cos \theta \frac{\partial}{\partial \theta} + i \sin \theta \frac{\partial}{\partial \varphi} \end{pmatrix}. \quad (25)$$

For the y component, the vectors are[7]

$$[\xi_{y,\hat{\mathbf{b}}}^+] = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\varphi/2} \\ ie^{i\varphi/2} \end{pmatrix} \quad \text{and} \quad [\xi_{y,\hat{\mathbf{b}}}^-] = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i\varphi/2} \\ -ie^{i\varphi/2} \end{pmatrix}. \quad (26)$$

However, it is found that in order to be able to derive a differential operator in this case, a phase factor $e^{-i\theta/2}$ must be introduced into the expressions for the eigenvectors. Thus, we have

$$[\xi_{y,\hat{\mathbf{b}}}^+] = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i(\varphi+\theta)/2} \\ ie^{i(\varphi-\theta)/2} \end{pmatrix} \quad \text{and} \quad [\xi_{y,\hat{\mathbf{b}}}^-] = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-i(\varphi+\theta)/2} \\ -ie^{i(\varphi-\theta)/2} \end{pmatrix}. \quad (27)$$

We find

$$[S_{y,\hat{\mathbf{b}}}] = \hbar \begin{pmatrix} -i \frac{\partial}{\partial \theta} & 0 \\ 0 & -i \frac{\partial}{\partial \theta} \end{pmatrix}. \quad (28)$$

The operator for the square of the spin is

$$[S^2] = [S_{x,\hat{\mathbf{b}}}]^2 + [S_{y,\hat{\mathbf{b}}}]^2 + [S_{z,\hat{\mathbf{b}}}]^2 = \hbar^2 \begin{pmatrix} i \frac{\partial}{\partial \varphi} - \frac{\partial^2}{\partial \varphi^2} & 0 \\ 0 & -i \frac{\partial}{\partial \varphi} - \frac{\partial^2}{\partial \varphi^2} \end{pmatrix} \quad (29)$$

The operators we have derived obey the usual commutation relations:

$$[[S_{i,\hat{\mathbf{b}}}], [S_{j,\hat{\mathbf{b}}}]] = i\hbar [S_{k,\hat{\mathbf{b}}}] \quad \text{with } i, j, k \text{ in cyclical order}, \quad (30)$$

and

$$[[S_{i,\hat{\mathbf{b}}}], [S^2]] = 0. \quad (31)$$

4 Conclusion

In this paper, we have shown that a differential eigenvalue equation for spin $1/2$ exists. We would hope to extend our arguments to other values of spin. Unlike the case of orbital angular momentum, it is clear that there is not one differential spin operator but as many as there are different values of spin. It would be surprising if differential eigenvalue equations did not exist for other values of spin. A very important question to be answered is the interpretation of the differential operator. This may conceivably lead to a clarification of the nature of intrinsic spin.

As the eigenfunctions satisfying our eigenvalue equation refer to the z direction as the initial direction of quantization, we have not obtained the most general form of the differential eigenvalue equation for spin $1/2$. Such a form would have an arbitrary direction as the initial direction of quantization. The search is proceeding for this general form.

5 References

1. Landé A., "From Dualism To Unity in Quantum Physics", Cambridge University Press, 1960.
2. Landé A., "New Foundations of Quantum Mechanics", Cambridge University Press, 1965.
3. Landé A., "Foundations of Quantum Theory," Yale University Press, 1955.
4. Landé A., "Quantum Mechanics in a New Key," Exposition Press, 1973.
5. Mweene H. V., "Derivation of Spin Vectors and Operators From First Principles", quant-ph/9905012
6. Mweene H. V., "New Treatment of Systems of Compounded Angular Momentum", quant-ph/9907082
7. Mweene H. V., "Generalized Spin-1/2 Operators and Their Eigenvectors", quant-ph/9906002